


List of Shrike-babblers found in India

Sno.	Name	Binomial Name
1	Black-eared Shrike-babbler	Pteruthius melanotis
2	Black-headed Shrike-babbler	Pteruthius rufiventer
3	Chestnut-fronted Shrike-babbler	Pteruthius aenobarbus aenobarbulus
4	Chestnut-winged Shrike-babbler	Pteruthius aeralatus validirostris
5	Green Shrike-babbler	Pteruthius xanthochlorus
6	Himalayan Shrike-babbler	Pteruthius flaviscapis ripleyi

Black-eared Shrike-babbler identification Tips

Black-eared Shrike-babbler: Pteruthius melanotis: Resident of Himalayas & hills of North East India



Black-headed Shrike-babbler identification Tips

Black-headed Shrike-babbler: Pteruthius rufiventer: Resident of Himalayas & hills of North East India

Iris is blue-grey to purplish-black, orbital skin grey Crown is glossy black with blue-green gloss Black lores Glossy black cheek and ear-coverts Upper mandible black to dusky, lower mandible pale grey or blue-grey Chestnut upperparts Chin and upper submoustachial area white Black wings blue-green gloss Pale grey on lower submoustachial area to mid-breast. Yellow patch on breast side Black tail with chestnut tips Mid-grey head side and neck side Yellow-tinged olive-green upperparts Black primary coverts, Brown-tinged pale pinkish underparts **Chestnut rump** Mid-grey Chin & throat Brown-tinged pale pinkish on belly Yellow-tinged olive-green Tail **Paler breast** Legs are pale pinkish-brown to chocolate-brown Male **Darker underparts Female Difference from male** Reference: Birds of Indian Subcontinent **Inskipp and Grimmett** ©www.ogaclicks.com Important id point

www.HBW.com


Chestnut-fronted Shrike-babbler identification Tips

Chestnut-fronted Shrike-babbler: Pteruthius aenobarbus aenobarbulus : Resident of Meghalaya in India

Iris is iris reddish to dark brown, White evering, broken front and rear Crown is yellow-tinged olive-green Dark chestnut lower forehead bordered above by short Greyish-white postocular supercilium bright yellow forecrown Yellow-green cheek and ear-coverts Yellow-tinged olive-green upperparts Bill is bluish-slate Chin and upper cheek to mid-throat dark chestnut Coverts black, broad white tips forming wing bars Chestnut breast olive-green Tail Olive brown upperparts Olive brown crown Yellow vent Wings with whitish fringes Wingbars pale rufous-buff Flanks are pale greyish-olive Yellow Belly & underparts Forehead is peach-buff Legs are pinkish **Paler Chin** Male Whitish from throat to underbelly Wings with greenish fringes **Difference from male** Reference: Birds of Indian Subcontinent **Inskipp and Grimmett** Important id point ©www.ogaclicks.com **Female** www.HBW.com


Chestnut-winged Shrike-babbler identification Tips

Chestnut-winged Shrike-babbler: Pteruthius aeralatus validirostris: Resident of hills of North East India



Green Shrike-babbler identification Tips

Green Shrike-babbler: Pteruthius xanthochlorus: Resident of Himalayas & hills of North East India




Difference from male
Important id point

Reference: Birds of Indian Subcontinent Inskipp and Grimmett www.HBW.com


Himalayan Shrike-babbler identification Tips

Himalayan Shrike-babbler: Pteruthius flaviscapis ripleyi : Resident of Himalayas




Comparison of Shrike-babbler - Male


Chestnut-winged Shrike-babbler



Himalayan Shrike-babbler



Black-headed Shrike-babbler



Comparison of Shrike-babbler - Female

